

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE May 31, 1996	3. REPORT TYPE AND DATES COVERED Final - 01/15/89 - 12/31/95		
4. TITLE AND SUBTITLE Final Technical Report of OCNR Grant N00014-89-J-1583 for the Support of Infrared Spatial Interferometry		5. FUNDING NUMBERS N00014-89-J-1583		
6. AUTHOR(S) Charles H. Townes				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) The Regents of the University of California c/o Sponsored Projects Office 336 Sproul Hall - 5940 University of California at Berkeley Berkeley, CA 94720-5940		8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) Office of Naval Research Attn: Lauranne Vogel Ballston Centre Tower One 800 North Quincy Street Arlington, VA 22217-5660		10. SPONSORING / MONITORING AGENCY REPORT NUMBER		
11. SUPPLEMENTARY NOTES		19960613 095		
12a. DISTRIBUTION / AVAILABILITY STATEMENT Unlimited		12b. DISTRIBUTION CODE		
13. ABSTRACT (Maximum 200 words) This grant has supported the development of infrared interferometry, studies of atmospheric transmission and image distortion, and the resolution and study of stars and material surrounding them, with the general goal of improving astrometry and hence determination of stellar positions. Substantial progress has been made, including determination of atmospheric distortions of infrared transmission and its variation with time, altitude location of major atmospheric distortions, correlation of radio and infrared measurements, and studies of possible change in apparent positions of stars due to stellar behavior. Findings are outlined briefly in this report. Rather complete discussion of the findings are in the 36 scientific publications supported by the grant. They are listed here and identified as to which of the several goals of the investigation they primarily apply.				
14. SUBJECT TERMS		15. NUMBER OF PAGES 8		
		16. PRICE CODE		
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT JL	

Final Technical Report of OCNR Grant N00014-89-J-1583
for the Support of Infrared Spatial Interferometry

Introduction

The grant N00014-89-J-1583 began on January 15, 1989 and ended December 31, 1995. Its purpose was the exploration and development of infrared spatial interferometry at wavelengths near 10 microns. The development had several goals. One was the general development of technology and exploration of performance of infrared interferometry. A second was study of fluctuations in pathlengths through the atmosphere and their characteristics, particularly as they affect the quality of interferometry or imaging through the atmosphere. A third was a coordination of measurements by radio interferometry of masers near stars with infrared interferometric measurements to determine where the masers occur with respect to the star, and with relevance to astrometry particularly in mind. A fourth was a general exploration of stellar characteristics with the infrared interferometer, with their relevance to astronomy particularly in mind. Progress in these various directions made during the term of the grant are discussed below.

Atmospheric Results

Fluctuations in the local index of refraction of the atmosphere have generally been modeled by the Kolmogorov approximation. This assumes an atmosphere of uniform average density, temperature, and humidity, and with no boundaries. This model predicts that the squares of differences between atmospheric indices of refraction at points separated by a distance R is proportional to $R^{2/3}$. It is also usually assumed that there is an upper limit to the distance between points for which the index refraction difference continues to increase. This upper limit is generally called the outer scale. There have been previously reported measurements indicating

that the Kolmogorov predictions are accurate representations of the actual behavior of the atmosphere and that the outer scale is very large, of the order of a kilometer. Results from the present investigation show that in fact the outer scale is often rather short, of the order of 10 meters or a few tens of meters, and that there are substantial deviations from the Kolmogorov predictions. The deviations are in the direction of making the build up of differences in index refraction between two points in the atmosphere somewhat slower than that predicted Kolmogorov model. All these results are quite favorable to imaging through the atmosphere and to interferometry. This is because previous expectations, based on the Kolmogorov approximation and on measurements in telescopes with rather limited size such as one or two meters diameter, would have predicted somewhat larger fluctuations than has been found here for large telescopes or interferometers with long baselines..

Our results are of course limited to the particular place of observation, namely the Mt. Wilson Observatory. However, measurements which have recently have been made at a number of other observatories indicate that an outer limit of the order of 10 meters or a few tens of meters is rather common and that in addition there are variations from the Kolmogorov increase of index refraction difference with distance in the same direction we have found on Mt. Wilson. How much the rate of increase of index of refraction differs is still under discussion, and this difference may be less at visible wavelengths than in the infrared. However, such differences do occur.

A second finding of the atmospheric measurements which is very encouraging for the future of imaging through the atmosphere is that a large fraction of the "poor seeing," or fluctuations in atmospheric pathlengths, is associated with a ground layer of the atmosphere not higher than about 40 meters. This occurs particularly when wind velocities are low, i.e., not higher than about 5 meters/second. On Mt. Wilson in particular, we find that about 2/3 of the disturbances in

pathlengths through the atmosphere come from a ground layer of atmosphere extending not higher than about 40 meters. These effects can fortunately be or corrected by local measurements of the lower level properties of the atmosphere, or they can be more or less eliminated by mounting telescopes high above the ground when this is practical.

Publications resulting from our work on atmospheric properties and pathlength fluctuations include numbers 8, 12, 13, 15, 16, 26, 27, and 33 of those listed at the end of this discussion.

Instrumentation

There has of course been a great deal of work on development and testing of the interferometer, including upgradings and modifications of the system which had been originally put together. Discussion of the instrument, its upgrading, and its performance are given in publications 2, 4, 10, 11, 18, 19, 20, 21, and 24 listed at the end of this discussion.

Astrometry

Although measurements of fluctuations in the atmosphere give a considerable amount of information of importance to astrometry, additional rather direct astrometric measurements have been made, including one particular case of a comparison between SiO masers as measured with radio interferometry and the radius of the dustshell around the star with which the masers were associated. This shows that SiO masers are in fact rather close to the surface of the star and where dust just begins to form. Discussions pertinent to these aspects of astrometry include publications 28, 29, 30, 31, and 35 in the list following this discussion.

Stellar Structure and Behavior

The size of the stars, formation of dust around them, distribution of this dust, and the changes of stars with time have been studied with the infrared interferometer. It appears that the sizes of stars can probably be better determined from infrared interferometry than from visible interferometry. It has been well known that stars vary in intensity or magnitude, and recently it has been found that some of this variation is associated with hot spots on the stars which change the center of the total radiation. We have shown that most of the variation in intensity of the bright star Betelgeuse is in fact associated with changes of temperature rather than size. These changes in temperature appear to be localized spots, and hence can change the center of gravity of the visible light emission, which affects high precision astrometry. The infrared is much less affected by such changes in temperature because it varies more or less linearly with temperature whereas visible light varies exponentially with temperature. Limb darkening due to the stellar atmosphere is also a less troublesome effect in the infrared than it is in the visible light, which is another reason why infrared measurements can give better determinations of the actual size of stars as well as their center of gravity.

We have found that many stars emit dust more or less episodically and irregularly. This dust is part of the reason their luminosity appears to change in the visible region, and can also cause some shift in the center of gravity of the visible light, which may affect astrometry.

In addition to the above comments concerning our study of stars, we have done extensive work on examining where the dust forms around stars, its distribution and temperature, and to some extent its composition. Such studies of stars are reported in publications 3, 5, 6, 7, 9, 14, 17, 23, 25, 32, 34, and 36 in the list following this discussion.

Collaborators

Students who have made significant contributions to work on this grant include David Hale, Peter McCullough, Everett Lipman, and John Monnier.

We have welcomed collaborators from other institutions who were interested in working with us on these technical and scientific problems. They include Dr. Thomas Geballe of the United Kingdom Infrared Telescope in Hawaii, who has measured certain infrared characteristics of the stars we have been studying in order to add to the detail with which interpretations of our results can be made. In addition, Dr. Lincoln Greenhill of the Smithsonian Astronomical Observatory in Cambridge, Massachusetts has collaborated with us in examining the relation between SiO masers about stars and the stellar characteristics. Dr. Robert Treuhaft of the Jet Propulsion Laboratory has collaborated with us in analyzing characteristics of the atmosphere, particularly the disturbed ground layer of the atmosphere, and the possibilities of accurate astrometry. Dr. Cuno Degiacomi of the University of Köln, Germany has collaborated with us in upgrading of the technology of the telescopes and in the measurement of stars. Dr. Bruno Lopez of the Observatory of Nice in France has also collaborated with us in examination of the variation of stars such as α Ceti with time and their deviation from sphericity.

Summary

Overall we believe that work which this grant has supported has made a very substantial impact on the technique of interferometry and particularly its use in the mid-infrared. We appreciate very much the important support provided by OCNR which has made this work possible.

Publications

1. "Infrared Long Baseline Interferometry," W.C. Danchi, M. Bester, P.R. McCullough, and C.H. Townes, *Highlights of Astronomy*, D. McNully (ed.) **8**, 563, D. Reidel, Dordrecht (1989).
2. "The U.C. Berkeley Infrared Spatial Interferometer: Instrumentation and First Results," M. Bester, W.C. Danchi, P.R. McCullough, and C.H. Townes, *Lecture Notes in Physics* **331**, 396, Springer, Heidelberg (1989).
3. "Spatially Resolved Dust Around IRC+10420," E.E. Bloemhoff, W.C. Danchi, and C.H. Townes, *Proceedings of the Sixth Cambridge Workshop on Cool Stars, Stellar Systems and the Sun*, (ASP) G. Wallerstein (ed.), **102**, 387 (1990).
4. "Long Baseline Interferometer for the Mid-infrared," M. Bester, W.C. Danchi, and C.H. Townes, *Amplitude and Intensity Spatial Interferometry*, J.B. Breckinridge (ed.), *Proc. SPIE* **1237**, 40 (1990).
5. "Visibility Curves at 10 μ m Wavelength for Stars and Dust Shells," W.C. Danchi, M. Bester, C. Degiacomi, and C.H. Townes, *Amplitude and Intensity Spatial Interferometry*, J.B. Breckinridge (ed.), *Proc. SPIE* **1237**, 327 (1990).
6. "Sub-arcsecond 10 micron Imaging of Circumstellar Dust," W.C. Danchi, *Astrophysics with Infrared Arrays*, R. Elston (ed.), *ASP Conference Series* **14**, 142 (1991).
7. "Location and Phase of Dust Formation in IRC+10216 Indicated by 11 μ m Spatial Interferometry," W.C. Danchi, M. Bester, C.G. Degiacomi, P.R. McCullough, and C.H. Townes *ApJ Lett* **359**, L59 (1990).
8. "Atmospheric Fluctuations in a 11 μ m Stellar Interferometer with Moveable Telescopes," M. Bester, C.G. Degiacomi, W.C. Danchi, and C.H. Townes, *Technical Digest of the Annual Meeting of the Optical Society of America*, Boston, MA, p. 82 (1990).
9. "Distribution of Dust About α Ceti and α Orionis Based on 11 μ m Spatial Interferometry," M. Bester, W.C. Danchi, C.G. Degiacomi, T.R. Geballe, and C.H. Townes *ApJ Lett* **367**, L27 (1991).
10. "Recent Astronomical Results from the Infrared Spatial Interferometer and Their Implications for a Lunar Optical/IR Interferometer," W.C. Danchi, M. Bester, P.R. McCullough, and C.H. Townes, *Proceedings of the Workshop on the Lunar Optical/IR Synthesis Array*, J.O. Burns (ed.) *NASA Conference Publication* **3066**, 75 (1992).
11. "Automation of Interferometric Observations at Mid-IR Wavelengths," M. Bester, C.G. Degiacomi, W.C. Danchi, L.J. Greenhill, and C.H. Townes, in *Robotic Telescopes in the 1990s*, A.V. Filippenko (ed.), *ASP Conference Series* **34**, 213 (1992).
12. "Atmospheric Pathlength Fluctuations in the Infrared Spatial Interferometer," C.H. Townes, *Adaptive Optics for Large Telescopes Technical Digest* (Optical Society of America, Washington, D.C.) **19**, 48 (1992).
13. "Atmospheric Fluctuations — Empirical Structure Functions and Projected Performance of Future Instruments," M. Bester, W.C. Danchi, C.G. Degiacomi, L.J. Greenhill, and C.H. Townes, *ApJ* **392**, 357 (1992).
14. "Short-exposure 'Speckle' Images of Dust Shells Around Late-type Stars," W.C. Danchi, C.G. Degiacomi, M. Bester, L.J. Greenhill, and D.Y. Gezari, in *High-Resolution Imaging by Interferometry II*, J.M. Beckers and F. Merkle (eds.), *ESO Conference and Workshop Proceedings* **39**, 21 (1992).
15. "Atmospheric Phase Fluctuations in the Infrared Spatial Interferometer," M. Bester, W.C. Danchi, C.G. Degiacomi, L.J. Greenhill, and C.H. Townes, in *Adaptive Optics for Large Telescopes Technical Digest* (Optical Society of America, Washington, DC) **19**, 48 (1992).

16. "Analysis of Atmospheric Fluctuations and Their Implications for High Resolution Astronomy," M. Bester, W.C. Danchi, C.G. Degiacomi, L.J. Greenhill, and C.H. Townes, in *High Resolution Imaging by Interferometry II*, J.M. Beckers and F. Merkle (eds.), *ESO Conference and Workshop Proceedings* **39**, 1017 (1992).
17. "Observations of Circumstellar Dust Shells with a Long Baseline Interferometer at 11 μm ," C.G. Degiacomi, M. Bester, W.C. Danchi, L.J. Greenhill, and C.H. Townes, in *High Resolution Imaging by Interferometry II*, J.M. Beckers and F. Merkle (eds.), *ESO Conference and Workshop Proceedings* **39**, 681 (1992).
18. "New Technical Developments at the Infrared Spatial Interferometer," C.G. Degiacomi, M. Bester, W.C. Danchi, L.J. Greenhill, and C.H. Townes, in *High Resolution Imaging by Interferometry II*, J.M. Beckers and F. Merkle (eds.), *ESO Conference and Workshop Proceedings* **39**, 721 (1992).
19. "Automation of Interferometric Observations," M. Bester, C. Degiacomi, W.C. Danchi, L. Greenhill, and C.H. Townes, *Robotic Telescopes in the 1990s*, *ASP Conference Series* **34**, 213 (1992).
20. "Recent Astronomical Results from the Infrared Spatial Interferometer and Their Implications for LOUISA," W.C. Danchi, M. Bester, and C.H. Townes, *Proc. of a Workshop on a Lunar Optical-Ultraviolet-Infrared Synthesis Array (LOUISA)*, J.O. Burns (ed.), *NASA Conference Publication* **3066**, 75 (1992).
21. "Heterodyne Interferometry in the Infrared," M. Bester, W.C. Danchi, C.G. Degiacomi, L.J. Greenhill, and C.H. Townes, *Very High Angular Resolution Imaging*, J.G. Robertson and W.J. Tango (eds.), **19**, 257, Kluwer (1993).
22. "Long Baseline Interferometric Observations of Circumstellar Dust Shells at 11 Microns," W.C. Danchi, L. Greenhill, M. Bester, C.G. Degiacomi, and C.H. Townes, *Very High Angular Resolution Imaging*, J.G. Robertson and W.J. Tango (eds.), **383**, Kluwer (1993).
23. "Characteristics of Dust Shells Around 13 Late-type Stars," W.C. Danchi, M. Bester, C.G. Degiacomi, L.J. Greenhill, and C.H. Townes, *AJ* **107**, 1469 (1994).
24. "Latest Technical Developments at the Infrared Spatial Interferometer," M. Bester, C.G. Degiacomi, W.C. Danchi, L.J. Greenhill, A. Riesinger, J. Weaver, and C.H. Townes, *Very High Angular Resolution Imaging*, J.G. Robertson and W.J. Tango (eds.), **19**, 257, Kluwer (1994).
25. "An IR Spatial Interferometer at 10 μm Wavelength and Measurement of Stellar Dust Shells," M. Bester, W.C. Danchi, C.G. Degiacomi, L. Greenhill, and C.H. Townes, *Infrared Physics and Technology*, F.K. Kneubühl (ed.) **35**, 503 (1994).
26. "The U.C. Berkeley Infrared Spatial Interferometer---Recent System Upgrades and Analysis of Atmospheric Fluctuations," M. Bester, W.C. Danchi, C.G. Degiacomi, and P.R. Bratt, *Amplitude and Intensity Interferometry II*, J.B. Breckinridge (ed.) *Proc. SPIE* **2200**, 274 (1994).
27. "Analysis of Visibility Data from the U.C. Berkeley Infrared Spatial Interferometer---Radiative Transfer Modeling and Results on Fifteen Stars," W.C. Danchi, M. Bester, L.J. Greenhill, C.G. Degiacomi, and C.H. Townes, *Amplitude and Intensity Spatial Interferometry II*, J.B. Breckinridge, (ed.), *Proc. SPIE* **2200**, 286 (1994).
28. "Toward Astrometric Tracking with the Infrared Spatial Interferometer," R.N. Treuhaft, M. Bester, W.C. Danchi, and C.H. Townes, *TDA Progress Report* **42-116**, 9 (1994).
29. "Circumstellar Envelope of VX Sgr Resolved with Long Baseline Interferometry at Millimeter and Mid-infrared Wavelengths," L.J. Greenhill, J.M. Moran, D.C. Becker, M. Bester, W.C. Danchi, and C.H. Townes, *Amplitude and Intensity Interferometry II*, J.B. Breckinridge (ed.) *Proc. SPIE* **2200**, 304 (1994).

30. "Interferometric Observations of the SiO Masers and Dust Shell of VX Sgr," L.J. Greenhill, F. Colomer, J.M. Moran, D.C. Backer, W.C. Danchi, and M. Bester, *ApJ* in press, 1995.
31. "Toward 10-Milliarcsecond Infrared Astrometry," R.N. Treuhaft, S.T. Lowe, M. Bester, W.C. Danchi, and C.H. Townes, *Amplitude and Intensity Spatial Interferometry II*, J.B. Breckinridge, (ed.) *Proc. SPIE* **2200**, 316 (1994).
32. "Long Baseline Interferometric Observations of Long Period Variable Stars," W.C. Danchi and M. Bester, to be published in *Proceedings of the Circumstellar Matter Conference*, G.D. Watt (ed.) *Astrophysics and Space Science* (1995).
33. "Vertical Scales of Turbulence at the Mt. Wilson Observatory," R.N. Treuhaft, S. Lowe, M. Bester, W.C. Danchi, and C.H. Townes, *ApJ* **453**, 522 (1995).
34. "Inner Radii of Dust Shells and Stellar Diameters Obtained by an Infrared Stellar Interferometer at 11 micron Wavelength," W.C. Danchi, M. Bester, L. Greenhill, C. Degiacomi, N. Geis, D. Hale, B. Lopez, and C.H. Townes, *Astrophysics and Space Science*, G. Watt and P. Williams (eds.), **224**, 447 (1995).
35. "The Circumstellar Envelope of VX Sgr," L.J. Greenhill, F. Colomer, J.M. Moran, M.J. Reid, D.C. Backer, M.A. Bester, and W.C. Danchi, *ApJ* **449**, 365 (1995).
36. "Measurement at 11 μ m Wavelengths of the Diameters of α Ori and α Sco; Changes in Effective Temperature of α Ori and Very Recent Dust Emission," M. Bester, W.C. Danchi, D. Hale, C.G. Degiacomi, D. Mekarnia, T.R. Geballe, and C.H. Townes, *ApJ* **463**, 336 (1996).